AOGS AS47: Interaction of Atmospheric Composition and the Asian Summer Monsoon

Validation of GOSAT SWIR xCH_4 using TCCON and Airborne Measurements

Chisa Iwasaki[1], Sachiko Hayashida[1], Toshinobu Machida[2], Andre Butz [3]

- Sachiko Hayashida < sachiko@ics.nara-wu.ac.jp > Nara Women's University, Japan
- Toshinobu Machida < tmachida@nies.go.jp > National Institute for Environmental Studies (NIES), Japan
- Andre Butz < andre.butz@kit.edu > IMK-ASF, Karlsruhe Institute of Technology (KIT), Germany

Abstract

As methane (CH₄) is one of the most important Short-Lived Climate Pollutants (SLCPs), global monitoring of atmospheric CH₄ with enough accuracy is expected to estimate its sources and sinks. For measurements of global distribution of CO₂ and CH₄ concentration from space, the Greenhouse gases Observing SATellite (GOSAT) was launched in 2009, and has continued measurements up to the present. However, though Monsoon Asia is one of the major source regions of CH₄, cloudy weather interfere satellite observation over the region. To understand CH₄ emission from this region, the selection of an adequate criterion of cloud screening, and validation of data quality are necessary.

In this study, we validate the GOSAT CH₄ products of the column-averaged dry-air mole fractions (xCH₄) derived from Short-Wavelength InfraRed (SWIR) radiation by comparing them with data of Total Carbon Column Observing Network (TCCON). Yoshida et al. (AMT, 2013) had already carried out the validation for the NIES product, but we extended the period and involved more TCCON sites; Yoshida et al. (2013) used data observed at the 13 TCCON sites from June, 2009 to December,

2012 and we used data at the 17 TCCON sites from June, 2009 to August, 2013. We found that the average difference between TCCON and GOSAT for the whole period is -6.0 ± 16.1 ppbv.

We also examined the appropriate cloud screening for xCH_4 product from RemoTeC-MACC. The product was obtained by using the "proxy method" which can provide us more data under cloudy conditions. In this study, we also tried to compare GOSAT data with aircraft measurements over Siberia and other areas.

Acknowledgements

This research was supported by the Environment Research and Technology Development Fund of the Ministry of the Environment, Japan (A1202).